Neuroscience Letters,184 (1995) 216–218
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abe, T., see Tohgi, H., 21
Akiguchi, I., see Nakamura, S., 29
Amadeo, A., Arcelli, P., Spreafico, R., De Biasi, S., Ultrastructural immunolocalization of muscarinic acetylcholine receptor in the dorsal thalamus of rat, 161
Anadón, R., see Carpintero, P., 63
Anand, P., Foley, P., Navsaria, H.A., Sinicropi, D., Williams-Chestnut, R.E., Leigh, I.M., Nerve growth factor levels in cultured human skin cells: effect of gestation and viral transformation, 157
Anderson, G.F., Clough-Helfman, C., Barraco, R.A., Displacement of norepinephrine by α-methylnorepinephrine in the nucleus tractus solitarius of the rat, 59
Arcelli, D., see Borasio, P.G., 97
Arcelli, P., see Amadeo, A., 161

Barraco, R.A., see Anderson, G.F., 59
Beyer, C., see Collado, P., 86
Boaten, A.S., see Boss, V., 1
Bocchini, V., see Murphy, Jr., G.M., 48
Boehm N., see Samama, B., 204
Borasio, P.G., Pavan, B., Fabbri, E., Ginanni-Corradini, F., Arcelli, D., Poli, A., Adenosine analogs inhibit acetylcholine release and cyclic AMP synthesis in the guinea-pig superior cervical ganglion, 97
Boss, V., Boaten, A.S., An L-cysteine sulfinic acid-sensitive metabotropic receptor mediates increased cAMP accumulation in hippocampal slices 1
Brett, R.R., see Pratt, J.A., 197

Carpintero, P., Anadón, R., del Amo, F.F., Gómez-Márquez, J., The thymosin β_4 gene is strongly activated in neural tissues during early postimplantation mouse development, 63
Castro, G.A., see Clatworthy, A.L., 5
Champagnat, J., see Pierrefiche, O., 101
Chan, S.H.H., see Shyr, M.-H., 212
Chateau, D., see Samama, B., 204
Chen, C.-F., see Shyr, M.-H., 212
Clatworthy, A.L., Illich, P.A., Castro, G.A., Walters, E.T., Role of periaxonal inflammation in the development of thermal hyperalgesia and guarding behavior in a rat model of neuropathic pain, 5
Clough-Helfman, C., see Anderson, G.F., 59
Coleman, H.A., see Holman, M.E., 75
Collado, P., Beyer, C., Hutchison, J.B., Holman, S.D., Hypothalamic distribution of astrocytes is gender-related in Mongolian gerbils, 86
Collet, A., see Pont, F., 52
Czurkó, A., see Fukuda, A., 13
Czyżewska-Szafran, H., see Jastrzębski, Z., 94

De Biasi, S., see Amadeo, A., 161
del Amo, F.F., see Carpintero, P., 63
Del Mar, N.A., see Geisert, Jr., E.E., 40
DeLanney, L.E., see Forno, L.S., 44
Deli, M.A., see Huszti, Z., 185

Eng, L.F., see Murphy, Jr., G.M., 48
Ernst, L.A., see Hossain, M.Z., 71
Escribá, P.V., see García-Sevilla, J.A., 133
Ewert, J.-P., see Spreckelsen, C., 105

Fabbri, E., see Borasio, P.G., 97
Fanardjian, V.V., Papoyan, E.V., Electrophysiological evidence for a direct neuronal connection from the motor cortex to the parietal association cortex of the cat, 201
Ferroni, S., see Mager, R., 165
Foley, P., see Anand, P., 157
Forno, L.S., DeLanney, L.E., Irwin, I., Langston, J.W., Ultrastructure of eosinophilic inclusion bodies in the amygdala-parahippocampal region of aged squirrel monkeys treated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, a dopaminergic neurotoxin, 44
Forsell, C., Lannfelt, L., Amyloid precursor protein mutation at codon 713 (Ala → Val) does not cause schizophrenia: non-pathogenic variant found at codon 705 (silent), 90
Fukuda, A., Czurkó, A., Hida, H., Muramatsu, K., Lénárd, L., Nishino, H., Appearance of deteriorated neurons on regionally different time tables in rat brain thin slices maintained in physiological condition, 13

García-Sevilla, J.A., Sastre, M., Escribá, P.V., Age-dependent increases of immunoreactive imidazoline receptors in the human brain: possible association of a 29/30 kDa protein with the I_2-imidazoline receptor identified by [^{3}H]idazoxan, 133
Geisert, Jr., E.E., Del Mar, N.A., Owens, J.L., Holmberg, E.G., Transfecting neurons and glia in the rat using pH-sensitive immunoliposomes, 40
Ginanni-Corradini, F., see Borasio, P.G., 97
Gómez-Márquez, J., see Carpintero, P., 63
Goźlińska, B., see Jastrzębski, Z., 94

Hamato, F., see Tohgi, H., 21
Hartung, H.-P., see Zielasek, J., 129
Hattori, A., Herbert, D.C., Vaughan, M.K., Yaga, K., Reiter, R.J., Melatonin inhibits luteinizing hormone releasing hormone (LHRH) induction of LH release from fetal rat pituitary cells, 109
Hawtin, S.R., see Pollack, S.J., 113
Heinemann, S.F., see Johnson, D.S., 67
Herbert, D.C., see Hattori, A., 109
Hida, H., see Fukuda, A., 13
Himi, T., see Sato, I., 145
Hirato, J., see Yokoo, H., 36
Holman, M.E., Tonta, M.A., Coleman, H.A., Parkington, H.C., Currents caused by the spontaneous release of quanta of acetylcholine onto chromaffin cells in guinea-pig adrenal medulla, 75
Holman, S.D., see Collado, P., 86
Holmberg, E.G., see Geisert, Jr., E.E., 40
Hossain, M.Z., Ernst, L.A., Nagy, J.I., Utility of intensely fluorescent cyanine dyes (CY3) for assay of gap junctional communication by dye-transfer, 71
Huszti, Z., Deli, M.A., Joó, F., Carrier-mediated uptake and release of histamine by cultured rat cerebral endothelial cells, 185
Hutchison, J.B., see Collado, P., 86

Illich, P.A., see Clatworthy, A.L., 5
Irwin, I., see Forno, L.S., 44

Jang, T.-H., see Tsai, C.-L., 208
Jastrzębski, Z., Czyżewska-Szafran, H., Goźlińska, B., Remiszewska, M., Mazurek, A.P., Clonidine hypotension in spontaneously hyper-

Neuroscience Letters,184 (1995) 216–218
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abe, T., see Tohgi, H., 21
Akiguchi, I., see Nakamura, S., 29
Amadeo, A., Arcelli, P., Spreafico, R., De Biasi, S., Ultrastructural immunolocalization of muscarinic acetylcholine receptor in the dorsal thalamus of rat, 161
Anadón, R., see Carpintero, P., 63
Anand, P., Foley, P., Navsaria, H.A., Sinicropi, D., Williams-Chestnut, R.E., Leigh, I.M., Nerve growth factor levels in cultured human skin cells: effect of gestation and viral transformation, 157
Anderson, G.F., Clough-Helfman, C., Barraco, R.A., Displacement of norepinephrine by α-methylnorepinephrine in the nucleus tractus solitarius of the rat, 59
Arcelli, D., see Borasio, P.G., 97
Arcelli, P., see Amadeo, A., 161

Barraco, R.A., see Anderson, G.F., 59
Beyer, C., see Collado, P., 86
Boaten, A.S., see Boss, V., 1
Bocchini, V., see Murphy, Jr., G.M., 48
Boehm N., see Samama, B., 204
Borasio, P.G., Pavan, B., Fabbri, E., Ginanni-Corradini, F., Arcelli, D., Poli, A., Adenosine analogs inhibit acetylcholine release and cyclic AMP synthesis in the guinea-pig superior cervical ganglion, 97
Boss, V., Boaten, A.S., An L-cysteine sulfinic acid-sensitive metabotropic receptor mediates increased cAMP accumulation in hippocampal slices 1
Brett, R.R., see Pratt, J.A., 197

Carpintero, P., Anadón, R., del Amo, F.F., Gómez-Márquez, J., The thymosin β_4 gene is strongly activated in neural tissues during early postimplantation mouse development, 63
Castro, G.A., see Clatworthy, A.L., 5
Champagnat, J., see Pierrefiche, O., 101
Chan, S.H.H., see Shyr, M.-H., 212
Chateau, D., see Samama, B., 204
Chen, C.-F., see Shyr, M.-H., 212
Clatworthy, A.L., Illich, P.A., Castro, G.A., Walters, E.T., Role of periaxonal inflammation in the development of thermal hyperalgesia and guarding behavior in a rat model of neuropathic pain, 5
Clough-Helfman, C., see Anderson, G.F., 59
Coleman, H.A., see Holman, M.E., 75
Collado, P., Beyer, C., Hutchison, J.B., Holman, S.D., Hypothalamic distribution of astrocytes is gender-related in Mongolian gerbils, 86
Collet, A., see Pont, F., 52
Czurkó, A., see Fukuda, A., 13
Czyżewska-Szafran, H., see Jastrzębski, Z., 94

De Biasi, S., see Amadeo, A., 161
del Amo, F.F., see Carpintero, P., 63
Del Mar, N.A., see Geisert, Jr., E.E., 40
DeLanney, L.E., see Forno, L.S., 44
Deli, M.A., see Huszti, Z., 185

Eng, L.F., see Murphy, Jr., G.M., 48
Ernst, L.A., see Hossain, M.Z., 71
Escribá, P.V., see García-Sevilla, J.A., 133
Ewert, J.-P., see Spreckelsen, C., 105

Fabbri, E., see Borasio, P.G., 97
Fanardjian, V.V., Papoyan, E.V., Electrophysiological evidence for a direct neuronal connection from the motor cortex to the parietal association cortex of the cat, 201
Ferroni, S., see Mager, R., 165
Foley, P., see Anand, P., 157
Forno, L.S., DeLanney, L.E., Irwin, I., Langston, J.W., Ultrastructure of eosinophilic inclusion bodies in the amygdala-parahippocampal region of aged squirrel monkeys treated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, a dopaminergic neurotoxin, 44
Forsell, C., Lannfelt, L., Amyloid precursor protein mutation at codon 713 (Ala → Val) does not cause schizophrenia: non-pathogenic variant found at codon 705 (silent), 90
Fukuda, A., Czurkó, A., Hida, H., Muramatsu, K., Lénárd, L., Nishino, H., Appearance of deteriorated neurons on regionally different time tables in rat brain thin slices maintained in physiological condition, 13

García-Sevilla, J.A., Sastre, M., Escribá, P.V., Age-dependent increases of immunoreactive imidazoline receptors in the human brain: possible association of a 29/30 kDa protein with the I_2-imidazoline receptor identified by [^{3}H]idazoxan, 133
Geisert, Jr., E.E., Del Mar, N.A., Owens, J.L., Holmberg, E.G., Transfecting neurons and glia in the rat using pH-sensitive immunoliposomes, 40
Ginanni-Corradini, F., see Borasio, P.G., 97
Gómez-Márquez, J., see Carpintero, P., 63
Goźlińska, B., see Jastrzębski, Z., 94

Hamato, F., see Tohgi, H., 21
Hartung, H.-P., see Zielasek, J., 129
Hattori, A., Herbert, D.C., Vaughan, M.K., Yaga, K., Reiter, R.J., Melatonin inhibits luteinizing hormone releasing hormone (LHRH) induction of LH release from fetal rat pituitary cells, 109
Hawtin, S.R., see Pollack, S.J., 113
Heinemann, S.F., see Johnson, D.S., 67
Herbert, D.C., see Hattori, A., 109
Hida, H., see Fukuda, A., 13
Himi, T., see Sato, I., 145
Hirato, J., see Yokoo, H., 36
Holman, M.E., Tonta, M.A., Coleman, H.A., Parkington, H.C., Currents caused by the spontaneous release of quanta of acetylcholine onto chromaffin cells in guinea-pig adrenal medulla, 75
Holman, S.D., see Collado, P., 86
Holmberg, E.G., see Geisert, Jr., E.E., 40
Hossain, M.Z., Ernst, L.A., Nagy, J.I., Utility of intensely fluorescent cyanine dyes (CY3) for assay of gap junctional communication by dye-transfer, 71
Huszti, Z., Deli, M.A., Joó, F., Carrier-mediated uptake and release of histamine by cultured rat cerebral endothelial cells, 185
Hutchison, J.B., see Collado, P., 86

Illich, P.A., see Clatworthy, A.L., 5
Irwin, I., see Forno, L.S., 44

Jang, T.-H., see Tsai, C.-L., 208
Jastrzębski, Z., Czyżewska-Szafran, H., Goźlińska, B., Remiszewska, M., Mazurek, A.P., Clonidine hypotension in spontaneously hyper-

tensive rats (SHR) depends on the functional state of GABAergic and glutamatergic systems, 94
Johnson, D.S., Heinemann, S.F., Detection of 5-HT_3R-A, a 5-HT_3 receptor subunit, in submucosal and myenteric ganglia of rat small intestine using in situ hybridization, 67
Joó, F., see Huszti, Z., 185
Jung, S., see Zielasek, J., 129

Kaneko, T., Minami, M., Satoh, M., Mizuno, N., Immunocytochemical localization of μ-opioid receptor in the rat caudate-putamen, 149
Kasahara, M., see Mizutani, T., 141
Kessey, K., see Sabo, S. 25
Kim, Y., see Sato, I., 145
Kimura, J., see Nakamura, S., 29
Kishi, H., see Tagoh, H., 17
Klein, W.L., see Sabo, S. 25
Knyihar-Csillik, E., Tajti, J., Mohtasham, S., Sari, G., Vecsei, L., Electrical stimulation of the Gasserian ganglion induces structural alterations of calcitonin gene-related peptide-immunoreactive perivascular sensory nerve terminals in the rat cerebral dura mater: a possible model of migraine headache, 189
Kolev, O.I., Milanov I., Vestibular and auditory influences on segmental motoneuron excitability – a comparative study, 193
Kovelowski, C.J., see Ossipov, M.H., 173
Krafft, G., see Sabo, S. 25
Kril, Y., see Mearow, K.M., 55
Kudo, M., see Takeo, T., 79
Künzig, H., see Zielasek, J., 129

Lai, Z., see Zhai, Q.Z., 82
Lallement, G., see Pont, F., 52
Lambert, M.P., see Sabo, S. 25
Langston, J.W., see Forno, L.S., 44
Lannfelt, L., see Forsell, C., 90
Lee, Y.L., see Murphy, Jr., G.M., 48
Leigh, I.M., see Anand, P., 157
Lénárd, L., see Fukuda, A., 13

Mager, R., Ferroni, S., Schubert, P., GTP- and GDP-analogues modulate an inwardly rectifying chloride channel in cultured hippocampal neurons, 165
Maki, B.E., see McIlroy, W.E., 177
Mazurek, A.P., see Jastrzębski, Z., 94
McIlroy, W.E., Maki, B.E., Early activation of arm muscles follows external perturbation of upright stance, 177
Mearow, K.M., Kril, Y., Anti-NGF treatment blocks the upregulation of NGF receptor mRNA expression associated with collateral sprouting of rat dorsal root ganglion neurons, 55
Mendelson, S.C., Quinn, J.P., Characterisation of potential regulatory elements within the rat preprotachykinin-A promoter, 125
Merlic, C.A., see Quinn, B., 169
Milanov I., see Kolev, O.I., 193
Minami, M., see Kaneko, T., 149
Mizuno, N., see Kaneko, T., 149
Mizutani, T., Kasahara, M., Degeneration of the intrahippocampal routes of the perforant and alvear pathways in senile dementia of Alzheimer type, 141
Mohtasham, S., see Knyihar-Csillik, E., 189
Motamed, S., see Quinn, B., 169
Muraguchi, A., see Tagoh, H., 17
Muramatsu, K., see Fukuda, A., 13
Murota, S.-i., see Sato, I., 145
Murphy, Jr., G.M., Song, Y., Ong, E., Lee, Y.L., Schmidt, K.G., Bocchini, V., Eng, L.F., Leukemia inhibitory factor mRNA is expressed in cortical astrocyte cultures but not in an immortalized microglial cell line, 48
Murthy, Ch.R.K., see Rama Rao, K.V., 137

Nagy, J.I., see Hossain, M.Z., 71
Nakamura, S., Akiguchi, I., Kimura, J., Topographic distributions of monoamine oxidase-B-containing neurons in the mouse striatum, 29
Nakazato, Y., see Yokoo, H., 36
Navsaria, H.A., see Anand, P., 157
Nishijo, H., see Tagoh, H., 17
Nishino, H., see Fukuda, A., 13
Nistri, A., see Stocca, G., 9
Nyberg, F., see Zhai, Q.Z., 82

Ong, E., see Murphy, Jr., G.M., 48
Ono, T., see Tagoh, H., 17
Ossipov, M.H., Kovelowski, C.J., Porreca, F., The increase in morphine antinociceptive potency produced by carrageenan-induced hindpaw inflammation is blocked by naltrindole, a selective δ-opioid antagonist, 173
Owens, J.L., see Geisert, Jr., E.E., 40

Papoyan, E.V., see Fanardjian, V.V., 201
Parkington, H.C., see Holman, M.E., 75
Pavan, B., see Borasio, P.G., 97
Pierrefiche, O., Champagnat, J., Richter, D.W., Calcium-dependent conductances control neurones involved in termination of inspiration in cats, 101
Poli, A., see Borasio, P.G., 97
Pollack, S.J., Sadler, I.I.J., Hawtin, S.R., Tailor, V.J., Shearman, M.S., Sulfated glycosaminoglycans and dyes attenuate the neurotoxic effects of β-amyloid in rat PC12 cells, 113
Pont, F., Collet, A., Lallement, G., Early and transient increase of rat hippocampal blood–brain barrier permeability to amino acids during kainic acid-induced seizures, 52
Porreca, F., see Ossipov, M.H., 173
Pratt, J.A., Brett, R.R., The benzodiazepine receptor inverse agonist FG 7142 induces cholecystokinin gene expression in rat brain, 197

Quinn, B., Toga, A.W., Motamed, S., Merlic, C.A., Fluoro Nissl Green: a novel fluorescent counterstain for neuroanatomy, 169
Quinn, J.P., see Mendelson, S.C., 125

Rama Rao, K.V., Vemuri, M.C., Murthy, Ch.R.K., Synaptosomal transport of branched chain amino acids in young, adult and aged rat brain cortex, 137
Redecker, P., The ras-like rab3A protein is present in pinealocytes of the gerbil pineal gland, 117
Reichmann, H., see Zielasek, J., 129
Reiter, R.J., see Hattori, A., 109
Remiszewska, M., see Jastrzębski, Z., 94
Richter, D.W., see Pierrefiche, O., 101
Roos, P., see Zhai, Q.Z., 82

Sabo, S., Lambert, M.P., Kessey, K., Wade, W., Krafft, G., Klein, W.L., Interaction of beta-amyloid peptides with integrins in a human nerve cell line, 25
Sadler, I.I.J., see Pollack, S.J., 113

Saheki, M., see Tohgi, H., 21
Sakuma, Y., see Takeo, T., 79
Samama, B., Chateau, D., Boehm N., Expression of NADPH-diaphorase in the rat forebrain during development, 204
Sari, G., see Knyihar-Csillik, E., 189
Sasaki, A., see Yokoo, H., 36
Sasaki, K., see Tohgi, H., 21
Sastre, M., see García-Sevilla, J.A., 133
Sato, I., Kim, Y., Himi, T., Murota, S.-i., Induction of calcium-independent nitric oxide synthase activity in cultured cerebellar granule neurons, 145
Satoh, M., see Kaneko, T., 149
Schmidt, K.G., see Murphy, Jr., G.M., 48
Schubert, P., see Mager, R., 165
Schürg-Pfeiffer, E., see Spreckelsen, C., 105
Shearman, M.S., see Pollack, S.J., 113
Shyr, M.-H., Tsai, T.-H., Tan, P.P.C., Chen, C.-F., Chan, S.H.H., Concentration and regional distribution of propofol in brain and spinal cord during propofol anesthesia in the rat, 212
Sinicropi, D., see Anand, P., 157
Song, Y., see Murphy, Jr., G.M., 48
Spreafico, R., see Amadeo, A., 161
Spreckelsen, C., Schürg-Pfeiffer, E., Ewert, J.-P., Responses of retinal and tectal neurons in non-paralyzed toads *Bufo bufo* and *B. marinus* to the real size versus angular size of objects moved at variable distance, 105
Stocca, G., Nistri, A., Enhancement of NMDA receptor mediated synaptic potentials of rat hippocampal neurones in vitro by thyrotropin releasing hormone, 9

Tagoh, H., Nishijo, H., Uwano, T., Kishi, H., Ono, T., Muraguchi, A., Reciprocal IL-1β gene expression in medial and lateral hypothalamic areas in SART-stressed mice, 17
Tailor, V.J., see Pollack, S.J., 113
Tajti, J., see Knyihar-Csillik, E., 189
Takahashi, S., see Tohgi, H., 21
Takeo, T., Kudo, M., Sakuma, Y., Stria terminalis conveys a facilitatory estrogen effect on female rat lordosis reflex, 79
Tan, P.P.C., see Shyr, M.-H., 212
Tatsuno, I., see Yamamoto, T., 32
Toga, A.W., see Quinn, B., 169
Tohgi, H., Abe, T., Saheki, M., Hamato, F., Sasaki, K., Takahashi, S., Reduced and oxidized forms of glutathione and α-tocopherol in the cerebrospinal fluid of parkinsonian patients: comparison between before and after L-dopa treatment, 21
Tomberg, C., New application of the Z estimator to identify the cognitive P300 in non-averaged human brain potentials, 153
Tonta, M.A., see Holman, M.E., 75
Toyka, , K.V., see Zielasek, J., 129
Tsai, C.-L., Jang, T.-H., Wang, L.-H., Effects of mercury on serotonin concentration in the brain of tilapia, *Oreochromis mossambicus* 208
Tsai, T.-H., see Shyr, M.-H., 212

Uwano, T., see Tagoh, H., 17

Vaughan, M.K., see Hattori, A., 109
Vecsei, L., see Knyihar-Csillik, E., 189
Vemuri, M.C., see Rama Rao, K.V., 137

Wade, W., see Sabo, S. 25
Walters, E.T., see Clatworthy, A.L., 5
Wang, H.-L., see Wu, T., 121
Wang, L.-H., see Tsai, C.-L., 208
Williams K., Pharmacological properties of recombinant *N*-methyl-D-aspartate (NMDA) receptors containing the ε4 (NR2D) subunit, 181
Williams-Chestnut, R.E., see Anand, P., 157
Wu, T., Wang, H.-L., Protein kinase C mediates neurotensin inhibition of inwardly rectifying potassium currents in rat substantia nigra dopaminergic neurons, 121

Yaga, K., see Hattori, A., 109
Yamamoto, T., Tatsuno, I., Antinociceptive effect of intrathecally administered pituitary adenylate cyclase activating polypeptide (PACAP) on the rat formalin test, 32
Yokoo, H., Sasaki, A., Hirato, J., Nakazato, Y., A neuronal fibrillary inclusion shares the epitope of p24 of human immunodeficiency virus, 36
Yukhananov, R., see Zhai, Q.Z., 82

Zhai, Q.Z., Lai, Z., Yukhananov, R., Roos, P., Nyberg, F., Decreased binding of growth hormone in the rat hypothalamus and choroid plexus following morphine treatment, 82
Zielasek, J., Reichmann, H., Künzig, H., Jung, S., Hartung, H.-P., Toyka, , K.V., Inhibition of brain macrophage/microglial respiratory chain enzyme activity in experimental autoimmune encephalomyelitis of the Lewis rat, 129